

Atlantic

Early Maincrop

(Wauseon x Lenape)



Consumer quality

Moderate yields of short oval tubers with high dry matter suitable for processing for crisps.



Agronomic features

Short dormancy with high dry matter ideal for crisp production in hot climates. Produces potatoes suitable for processing in 90 days. Resistant to PCN Ro1 and immune to PVX.

Tuber characteristics

| | | | |
|----------------------|----------|--------------------|----|
| Market Sector | C | Shape of tuber | SO |
| Maturity | 3 | Depth of eyes | 3 |
| Colour of skin | C | Smoothness of skin | M |
| Colour of flesh | C | | |

Resistance to damage, pests and diseases

| | | | |
|---------------------|---|----------------------------------|---|
| Foliage blight | 3 | Virus Y | 3 |
| Tuber blight | 3 | Skin Spot | – |
| Blackleg | 5 | Dry Rot (<i>F. coeruleum</i>) | – |
| Common scab | 6 | Dry Rot (<i>F. sulphureum</i>) | – |
| Powdery scab | 2 | Silver Scurf | – |
| G.Pallida Pa 2/3,1 | 2 | Bruising | 3 |
| G.Rostochiensis Ro1 | 9 | Splitting | 3 |
| Leafroll | 6 | | |